

# It's critical TO VACCINATE

to ensure pets have a great stay

## The most cost-effective, safest way to prevent disease<sup>1-4</sup>

### NEW CLIENTS

Review vaccine history prior to stay (clients can request records from their vet clinic)

### RETURNING CLIENTS

Ensure no recent history of vomiting, diarrhea, sneezing, coughing, skin infections, fever, or new illnesses

### ALL CLIENTS

Confirm no new vaccines will be needed during the stay-if so, vaccinations must be completed BEFORE they arrive

- **Canine combo:** 2-dose initial series or booster 3 weeks before stay
- **CIV (dog flu):** 2-dose initial series or booster 2 weeks before stay
- **Bordetella (canine cough):** 7 days before stay

## KEY VACCINES FOR DOGS AND CATS<sup>1,2</sup>

### DOGS



**Canine Combo (DAPP):** Every 3 years after initial series

- Canine Distemper Virus
- Canine Adenovirus-1 & 2
- Canine Parainfluenza Virus
- Canine Parvovirus

**Rabies\*:** Every 1 or 3 years, depending on local laws

**Leptospirosis (L4)\*:** Yearly after initial series

- Given alone or in combination with DAPP
- Covers 4 subtypes of leptospirosis

**Bordetella bronchiseptica (canine cough)\*:** Yearly after initial series


- May be given in combination with Canine Parainfluenza and Canine Adenovirus boosters based on local risk

**Canine Influenza Virus (CIV, or dog flu):** Yearly after initial series

- Bivalent CIV H3N2/H3N8 covers both known strains

\*This disease may infect humans.<sup>5</sup>

### CATS



**Feline Combo (RCP):** Every 3 years after initial series

- Feline Viral Rhinotrachitis
- Feline Calicivirus
- Feline Panleukopenia

**Rabies\*:** Every 1 or 3 years, depending on local laws

**Feline Leukemia Virus (FeLV):** Yearly after initial series

**DISEASE CAN SPREAD FROM ANIMALS TO SURFACES TO OTHER ANIMALS AND PEOPLE<sup>3</sup>**



Coughing, sneezing



Contact between animals



Urine, diarrhea, vomiting



Food/water bowls, toys

**ANYWHERE PETS ARE TOGETHER THEY ARE AT GREATEST RISK FOR DISEASE OUTBREAKS**

What we do to keep pets healthy<sup>3</sup>



Require vaccinations



Cleaning/disinfection



Handwashing



Strict use of work only clothing



Rapid isolation of sick animals

References: 1. Ford RB, Larson LJ, McClure KD, Schultz RD, Welborn LV. 2017 AAHA canine vaccination guidelines. Trends. October 2017;26-35. 2. Scherk MA, Ford RB, Gaskell RM, et al. 2013 AAFP feline vaccination advisory panel report. J Feline Med Surg. 2013;15:785-808. 3. Stull JW, Sherding RG, O'Quin J, et al. Infectious Disease in Dogs in Group Settings: Strategies to Prevent Infectious Diseases in Dogs at Dog Shows, Sporting Events, and Other Canine Group Settings. September 2016. 4. Stull JW, Kasten JI, Evason MD, et al. Risk reduction and management strategies to prevent transmission of infectious disease among dogs at dog shows, sporting events, and other canine group settings. J Am Vet Med Assoc. 2016;249(6):612-627. 5. Ghasemzadeh I, Namazi SH. Review of bacterial and viral zoonotic infections transmitted by dogs. J Med Life. 2015;8(4):1-5.

PROVIDED AS A SERVICE OF

All trademarks are the property of their respective owners.  
© 2019 Zoetis Services LLC. All rights reserved. SAB-00814 B

**ZOETIS  
PETCARE**

**VANGUARD**

**RESCUE  
DISINFECTANTS**

**VIROX™  
ANIMAL HEALTH**